



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Exanthema associated with *Capnocytophaga canimorsus* bacteremia

Uçkay, Ilker ; Stirnemann, Jérôme

DOI: <https://doi.org/10.1016/j.ijid.2019.03.006>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-170097>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Uçkay, Ilker; Stirnemann, Jérôme (2019). Exanthema associated with *Capnocytophaga canimorsus* bacteremia. *International Journal of Infectious Diseases*, 82:104-105.

DOI: <https://doi.org/10.1016/j.ijid.2019.03.006>



Medical Imagery

Exanthema associated with *Capnocytophaga canimorsus* bacteremia

ARTICLE INFO

Article history:

Received 27 November 2018

Received in revised form 4 March 2019

Accepted 6 March 2019

Corresponding Editor: Eskild Petersen, Aarhus, Denmark

Keywords:
Exanthema
Capnocytophaga
Sepsis

A 51-year-old healthy sportsman was admitted with a history of transient fever and non-itchy exanthema all over his body, sparing the oral mucosa (Figure 1) (Jordan et al., 2016). His haemodynamic parameters were normal. The patient denied a history of chills, unusual sexual contact, trauma, or tick bites, but reported a recent forest walk and swimming in the river with his dog. The dog, which had remained in good health, had probably caused him a minor injury. With a clinical suspicion of borreliosis (first serology negative), we administered oral doxycycline 100 mg twice daily empirically. The next day he left hospital and we received the results of the blood cultures: *Capnocytophaga canimorsus*. Due to his excellent clinical progress, we continued doxycycline and added amoxicillin–clavulanic acid without switching to intravenous therapy, both for another 3 weeks. There were two reasons for

adding another oral antimicrobial drug. The first was the uncertainty regarding the clinical efficacy of oral doxycycline for *Capnocytophaga* bacteremia. Although aminopenicillins are the usual recommended first-line agents for *C. canimorsus* (Jolivet-Gougeon et al., 2007; Sotiriou et al., 2015), numerous reports have stated its susceptibility to doxycycline (Jolivet-Gougeon et al., 2007) and to many other drugs, while most strains remain resistant to polymyxins, fosfomycin, and trimethoprim (Jolivet-Gougeon et al., 2007). The second reason was the limited experience of oral doxycycline for the treatment of bacteremia that has been reported so far in combination with gentamicin for *Bartonella* or *Brucella* spp (Fruchtman et al., 2015). The exanthema, which was different from the usual livedo racemosa seen in *C. canimorsus* sepsis (Sotiriou et al., 2015), disappeared and the



Figure 1. Non-itchy exanthema all over the patient's body, sparing the oral mucosa.

control serology for *Borrelia* remained negative. The patient has reported being in good health 1.5 years after the infection. We share the photos of the patient's exanthema associated with *C. canimorsus* bacteremia (Jordan et al., 2016).

Funding

No funding for this article.

Patient consent

The patient consented by signature to the publication of his history and photographs.

Conflict of interest

The authors have no conflict of interest to declare. This publication fulfils the ethical requirements of the Declaration of Helsinki.

References

Fruchtman Y, Segev RW, Golan AA, Dalem Y, Tailakh MA, Novak V, et al. Epidemiological, diagnostic, clinical, and therapeutic aspects of *Brucella* bacteremia in children in southern Israel: a 7-year retrospective study (2005–2011). *Vector Borne Zoonotic Dis* 2015;15:195–201.

Jolivet-Gougeon A, Sixou JL, Tamanai-Shacoori Z, Bonnaure-Mallet M. Antimicrobial treatment of *Capnocytophaga* infections. *Int J Antimicrob Agents* 2007;29:367–73.

Jordan CS, Minitier U, Yarbrough K, Mengden SJ. Urticarial exanthem associated with *Capnocytophaga canimorsus* bacteremia after a dog bite. *JAAD Case Rep* 2016;2:98–101.

Sotiriou A, Sventzouri S, Nepka M, Magira EE. Acute generalized livedo racemosa caused by *Capnocytophaga canimorsus* identified by MALDI-TOF MS. *Int J Infect Dis* 2015;33:196–8.

Ilker Uçkay^{a,*}

Jérôme Stirnemann^b

^aBalgrist University Hospital, Zurich, Switzerland

^bDepartment of Internal Medicine, Geneva University Hospitals and Medical School, Geneva, Switzerland

* Corresponding author at: Balgrist University Hospital, Forchstrasse 340, 8008 Zurich, Switzerland.

E-mail address: ilker.uckay@balgrist.ch (I. Uçkay).

Corresponding Editor: Eskild Petersen, Aarhus, Denmark

Received 27 November 2018

Received in revised form 4 March 2019

Accepted 6 March 2019